

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

### Abstract of the Disclosure

Methods and compositions useful for inducing a cytotoxic T lymphocyte response (CTL) in a human or domesticated or agriculturally important animal. The method includes the steps of providing the antigen to which the CTL response is desired and providing a microfluidized antigen formulation which comprises, consists, or consists essentially of two or more of a stabilizing detergent, a micelle-forming agent, and an oil. This antigen formulation is preferably lacking in an immunostimulating peptide component, or has sufficiently low levels of such a component that the desired CTL response is not diminished. This formulation is provided as a stable oil-in-water emulsion.